

On the maximal order of numbers in the “factorisatio numerorum” problem

Martin Klazar

Department of Applied Mathematics (KAM)
and Institute for Theoretical Computer Science (ITI), Charles University
Malostranské náměstí 25, 118 00 Praha, Czech Republic
`klazar@kam.mff.cuni.cz`

Florian Luca

Instituto de Matemáticas, Universidad Nacional Autónoma de México
Ap. Postal 61-3 (Xangari), C.P. 58089, Morelia, Michoacán, México
`fluca@matmor.unam.mx`

February 1, 2008

Abstract

Let $m(n)$ be the number of ordered factorizations of $n \geq 1$ in factors larger than 1. We prove that for every $\varepsilon > 0$

$$m(n) < \frac{n^\rho}{\exp((\log n)^{1/\rho}/(\log \log n)^{1+\varepsilon})}$$

holds for all integers $n > n_0$, while, for a constant $c > 0$,

$$m(n) > \frac{n^\rho}{\exp(c(\log n/\log \log n)^{1/\rho})}$$

holds for infinitely many positive integers n , where $\rho = 1.72864\dots$ is the real solution to $\zeta(\rho) = 2$. We investigate also arithmetic properties of $m(n)$ and the number of distinct values of $m(n)$.

1 Introduction

Let $m(n)$ be the number of ordered factorizations of a positive integer n in factors bigger than 1. For example, $m(12) = 8$ since we have the factorizations 12 , $2 \cdot 6$, $6 \cdot 2$, $3 \cdot 4$, $4 \cdot 3$, $2 \cdot 2 \cdot 3$, $2 \cdot 3 \cdot 2$, and $3 \cdot 2 \cdot 2$. By the definition, $m(1) = 0$ but we will see that in some situations it is useful to set $m(1) = 1$ or $m(1) = 1/2$. Kalmár [13] found the average order of $m(n)$: for $x \rightarrow \infty$,

$$M(x) = \sum_{n \leq x} m(n) = cx^\rho(1 + o(1)), \quad (1)$$

where $\rho = 1.72864\dots$ is the real solution to $\zeta(\rho) = 2$ and $c = 0.31817\dots$ is given by $c = -1/\rho\zeta'(\rho)$. (As usual, $\zeta(s) = \sum_{n \geq 1} n^{-s}$.) Further detailed and strong results on the average order of $m(n)$ were obtained by Hwang [9].

In contrast, good bounds on the maximal order of $m(n)$ were lacking. Erdős claimed in the end of his article [4] that there exist positive constants $0 < c_2 < c_1 < 1$ such that

$$m(n) < \frac{n^\rho}{\exp((\log n)^{c_2})}$$

holds for all $n > n_0$, while

$$m(n) > \frac{n^\rho}{\exp((\log n)^{c_1})}$$

holds for infinitely many n , but he gave no details. To our knowledge, the best proved bounds on the maximal order state that $m(n) < n^\rho$ for every $n \geq 1$ (Chor, Lemke and Mador [1], a simple proof by induction was recently given by Coppersmith and Lewenstein [3]) and that for any $\varepsilon > 0$ one has $m(n) > n^{\rho-\varepsilon}$ for infinitely many n (Hille [8], [3] gives an explicit construction). (In Lemma 2.4 we strengthen the argument of [1] and show that $m(n) \leq n^\rho/\sqrt{2}$ for every $n \geq 1$.)

Here we come close to determining the maximal order of $m(n)$. We prove that it is, roughly, $n^\rho/\exp((\log n)^{1/\rho})$. More precisely, we prove that for every $\varepsilon > 0$,

$$m(n) < \frac{n^\rho}{\exp((\log n)^{1/\rho}/(\log \log n)^{1+\varepsilon})}$$

holds for all $n > n_0$ (Theorem 3.1), while

$$m(n) > \frac{n^\rho}{\exp(c(\log n)^{1/\rho}/(\log \log n)^{1/\rho})}$$

holds with certain constant $c > 0$ for infinitely many positive integers n (Theorem 4.1).

The paper is organized as follows. In Section 2 we give auxiliary results, of which Lemma 2.3 on the speed of convergence $\rho_k \rightarrow \rho$ (ρ_k is a “finite” counterpart of ρ for $m(n)$ restricted to smooth numbers n with no prime factor exceeding p_k , the k th prime number) and Lemmas 2.4–2.6 giving explicit inequalities for $m(n)$ and $m_k(n)$ ($m_k(n) = m(n)$ if n has no prime factor $> p_k$ and $m_k(n) = 0$ else) may be of independent interest. Section 3 is devoted to the proof of the upper bound. The proof is elementary (uses real analysis only) and is obtained by combining the combinatorial bounds on $m(n)$ in Lemmas 2.4 and 2.5, standard bounds from the prime numbers theory, and the convergence bound in Lemma 2.3. Section 4 is devoted to the proof of the lower bound. In the first version of this article, still available at [15, version 1], we proved by an elementary approach similar to that in Section 3, with the additional ingredient being Kalmár’s asymptotic relation (1), a weaker lower bound that has $(\log n)^{1/\rho}$ in the denominator replaced with the bigger power $(\log n)^{\rho/(\rho^2-1)+o(1)}$. Here, we prove in Section 4 a lower bound with the matching exponent $1/\rho$ of the $\log n$ by a method suggested to us by an anonymous referee. The method works in the complex domain and combines the uniform (i.e., with error estimates independent on k) version of (1) for $m_k(n)$, bounds on smooth numbers, and again Lemma 2.3. In Section 5, we give further references and comments on the history of $m(n)$ and some related problems. We also investigate arithmetical properties of $m(n)$ and prove, for example, that $m(n)$ is not eventually periodic modulo k for any integer $k > 1$, and that $m(n)$ is not a holonomic sequence.

Acknowledgments. Most of this paper was written during a very enjoyable visit by the first author to the Mathematical Institute of the UNAM in Morelia, Mexico, in March 2005. This author wishes to express his thanks to that institution for the hospitality and support. He also acknowledges the support to ITI by the project 1M0021620808 of the Czech Ministry of Education. Both authors are deeply grateful to an anonymous referee who outlined for them a plan how to prove a better lower bound, which is now carried out in Section 4.

2 Preliminaries and auxiliary results

Let us begin with recalling some notation. For a positive integer n we write $\omega(n)$ and $\Omega(n)$ for the number of distinct prime factors of n and the total number of prime factors of n (including multiplicities), respectively. We use the letters p and q with or without subscripts to denote prime numbers. We put $P(n)$ for the largest prime factor of n . We write $\log$ for the natural logarithm. In the complex domain (mainly in Section 4) we use s to denote generic variable and write σ and τ for its real and imaginary part, respectively, so $s = \sigma + i\tau$, where $i = \sqrt{-1}$. We use the Vinogradov symbols $\ll$ and $\gg$ and the Landau symbols O and o with their usual meanings.

The proof of the following estimate is standard and we omit it.

Lemma 2.1. *If $\delta > \delta_0 > 1$, then the estimate*

$$\sum_{p>t} \frac{1}{p^\delta} = \frac{(\delta-1)^{-1}}{t^{\delta-1} \log t} + O\left(\frac{1}{t^{\delta-1}(\log t)^2}\right) \quad (2)$$

holds uniformly for $t > 2$.

Let p_k be the k th prime. We shall use the well known asymptotic relations

$$\sum_{p \leq x} \log p = x + O(x/\log x)$$

(equivalent to the Prime Number Theorem) and

$$p_k = k \log k + k \log \log k + O(k)$$

(the full asymptotic expansion $p_k = k(\log k + \log \log k - 1 + \dots)$ was found by Cipolla [2]). Let $\mathcal{P}_k$ be the set (including 1) of positive integers composed only of the primes $p_1 = 2, p_2, \dots, p_k$, and $m_k(n)$ be the number of ordered factorizations of n in factors lying in $\mathcal{P}_k \setminus \{1\}$. We allow $k = \infty$, then $p_k = \infty$, $\mathcal{P}_\infty = \mathcal{P}$ is the set of all primes, and $m_\infty(n) = m(n)$. Note that, for $k \in \mathbb{N}$, $m_k(n) > 0$ iff $n \in \mathcal{P}_k$, if $m_k(n) > 0$ then $m_k(n) = m(n)$, and if $n \leq p_k$ then $m_k(n) = m(n)$. Let, for complex s with $\sigma > 1$ and $k \in \mathbb{N} \cup \{\infty\}$,

$$\zeta_k(s) = \prod_{p \leq p_k} \left(1 - \frac{1}{p^s}\right)^{-1} = \sum_{n \in \mathcal{P}_k} \frac{1}{n^s}$$

and ρ_k be the real solution to $\zeta_k(\rho_k) = 2$. For $k = \infty$ we get the Euler-Riemann zeta function $\zeta(s) = \zeta_\infty(s)$ and the number $\rho = \rho_\infty$. Note that for $k \in \mathbb{N}$ the series for $\zeta_k(s)$ converges absolutely even for $\sigma > 0$. For every s with $\sigma > 1$ we have the convergence $\zeta_k(s) \rightarrow \zeta(s)$ as $k \rightarrow \infty$. For $k \in \mathbb{N} \cup \{\infty\}$, one has the identity (setting $m_k(1) = 1$ for every k)

$$\sum_{n \geq 1} \frac{m_k(n)}{n^s} = \sum_{l \geq 0} (\zeta_k(s) - 1)^l = \frac{1}{2 - \zeta_k(s)},$$

which implies that $m_k(n) = o(n^{\rho_k + \varepsilon})$ for every fixed $\varepsilon > 0$. Our approach to estimating $m(n)$ is based on approximating the “infinite” quantities $m(n)$, ρ , and $\zeta(s)$ with their “finite” counterparts $m_k(n)$, ρ_k , and $\zeta_k(s)$ for $k \in \mathbb{N}$ but $k \rightarrow \infty$. We quantify the degrees of approximation in the following two lemmas. The first lemma is obtained by considering the infinite series defining $\zeta_k(s)$ and $\zeta(s)$ and its easy proof is omitted.

Lemma 2.2. *We have*

$$\rho_1 = 1 < \rho_2 = 1.43527 \dots < \rho_3 = 1.56603 \dots < \dots < \rho = 1.72864 \dots$$

and $\rho_k \rightarrow \rho$ as $k \rightarrow \infty$. The convergence $\zeta_k(s) \rightarrow \zeta(s)$ as $k \rightarrow \infty$ is uniform on every complex domain $\sigma > \sigma_0 > 1$ and the same is true for the convergence $\zeta'_k(s) \rightarrow \zeta'(s)$ and for all higher derivatives. Also, for every $k \in \mathbb{N} \cup \{\infty\}$ we have $\zeta'_k(\rho_k) < 0$.

We shall use this lemma to bound various expressions containing ρ_k , $\zeta_k(\rho_k)$, $\zeta_k(s)$, $1/\zeta'_k(\rho_k)$ etc. by constants independent on k .

Lemma 2.3. *The estimate*

$$\rho - \rho_k = \frac{1}{(\rho - 1)|\zeta'(\rho)|} \cdot \frac{1}{k^{\rho-1}(\log k)^\rho} \left(1 + O\left(\frac{\log \log k}{\log k}\right) \right)$$

holds for all $k \geq 2$.

Proof. We will assume that $k \geq 2$. The equation $\zeta_k(\rho_k)^{-1} = \zeta(\rho)^{-1} = 1/2$ implies that

$$\prod_{2 \leq p \leq \rho_k} \left(1 - \frac{1}{p^{\rho_k}} \right) = \prod_{p \geq 2} \left(1 - \frac{1}{p^\rho} \right).$$

Taking logarithms and regrouping, we get

$$\sum_{2 \leq p \leq p_k} \left(\log \left(1 - \frac{1}{p^\rho} \right) - \log \left(1 - \frac{1}{p^{\rho_k}} \right) \right) = - \sum_{p > p_k} \log \left(1 - \frac{1}{p^\rho} \right).$$

The left side satisfies, by Lagrange's Mean-Value Theorem (the derivative of the function $x \mapsto \log(1 - 1/p^x)$ is $(\log p)/(p^x - 1)$),

$$\begin{aligned} \sum_{2 \leq p \leq p_k} \log \left(1 - \frac{1}{p^\rho} \right) - \log \left(1 - \frac{1}{p^{\rho_k}} \right) &= (\rho - \rho_k) \sum_{2 \leq p \leq p_k} \frac{\log p}{p^{\sigma_p} - 1} \quad (3) \\ &> (\rho - \rho_k)(\log 2)/3 \end{aligned}$$

for some numbers $\sigma_p \in (\rho_k, \rho) \subset (1.4, 1.8)$. The right side is

$$\begin{aligned} - \sum_{p > p_k} \log \left(1 - \frac{1}{p^\rho} \right) &= \sum_{p > p_k} \frac{1}{p^\rho} + O \left(\sum_{p > p_k} \frac{1}{p^{2\rho}} \right) \\ &= \frac{(\rho - 1)^{-1}}{p_k^{\rho-1} \log(p_k)} \left(1 + O \left(\frac{1}{\log k} \right) \right) \\ &= \frac{(\rho - 1)^{-1}}{k^{\rho-1} (\log k)^\rho} \left(1 + O \left(\frac{\log \log k}{\log k} \right) \right), \quad (4) \end{aligned}$$

where we used Lemma 2.1 and the fact that $p_k = k(\log k + O(\log \log k))$. We get immediately that

$$\rho - \rho_k \ll \frac{1}{k^{\rho-1} (\log k)^\rho}. \quad (5)$$

To do better, we return to (3) and write

$$\frac{\log p}{p^{\sigma_p} - 1} = \frac{\log p}{p^\rho - 1} \left(1 + \frac{p^{\sigma_p}}{p^{\sigma_p} - 1} (p^{\rho - \sigma_p} - 1) \right).$$

We have $1 \leq p^{\sigma_p}/(p^{\sigma_p} - 1) \leq 2$ and, using (5),

$$p^{\rho - \sigma_p} - 1 \leq \exp((\rho - \rho_k) \log p_k) - 1 \ll (\rho - \rho_k) \log p_k \ll \frac{1}{k^{\rho-1} (\log k)^{\rho-1}}.$$

Hence the right side of (3) equals

$$\begin{aligned} (\rho - \rho_k) \sum_{2 \leq p \leq p_k} \frac{\log p}{p^{\sigma_p} - 1} &= (\rho - \rho_k) (1 + O(k^{1-\rho} (\log k)^{1-\rho})) \sum_{2 \leq p \leq p_k} \frac{\log p}{p^\rho - 1} \\ &= (\rho - \rho_k) (1 + O(k^{-1/2})) \sum_{2 \leq p \leq p_k} \frac{\log p}{p^\rho - 1}. \end{aligned}$$

Equating the right side of (3) and (4) we get the relation

$$(\rho - \rho_k) \sum_{2 \leq p \leq p_k} \frac{\log p}{p^\rho - 1} = \frac{(\rho - 1)^{-1}}{k^{\rho-1}(\log k)^\rho} \left(1 + O\left(\frac{\log \log k}{\log k}\right) \right).$$

All is left to notice is that

$$\begin{aligned} |\zeta'(\rho)| &= \sum_{p \geq 2} \frac{\log p}{p^\rho - 1} = \sum_{p \leq p_k} \frac{\log p}{p^\rho - 1} + \sum_{p > p_k} \frac{\log p}{p^\rho - 1} \\ &= \sum_{p \leq p_k} \frac{\log p}{p^\rho - 1} + O(k^{-1/2}), \end{aligned}$$

where the last estimate follows again from Lemma 2.1 via the fact that $\log p \ll p^{1/10}$:

$$\sum_{p > p_k} \frac{\log p}{p^\rho - 1} \ll \sum_{p > p_k} \frac{1}{p^{\rho-0.1}} \ll \frac{1}{p_k^{\rho-1.1} \log p_k} < 1/\sqrt{k}.$$

The claimed estimate now follows. $\square$

In the next three lemmas, we prove combinatorial inequalities involving $m_k(n)$ and $m(n)$. In the first lemma, we slightly improve the result from [1, Theorem 5] that $m_k(n) < n^{\rho_k}$ for every $n \geq 1$. The second lemma is crucial for obtaining bounds of the type $m(n) = o(n^\rho)$. The third lemma gives some lower estimates on $m(n)$.

Lemma 2.4. *For every $k \in \mathbb{N} \cup \{\infty\}$ and $n \geq 1$ (with $m_k(1) = 0$),*

$$m_k(n) \leq \frac{1}{\sqrt{2}} n^{\rho_k}.$$

Proof. For every $r, s \geq 1$ we have (now setting $m_k(1) = 0$)

$$m_k(rs) \geq 2m_k(r)m_k(s). \tag{6}$$

To show this inequality, we assume that $r, s \geq 2$ (for $r = 1$ or $s = 1$ it holds trivially) and consider the set X of all pairs (u, v) where u (v) is an ordered factorization of r (s) in factors lying in $\mathcal{P}_k \setminus \{1\}$, and the set Y of the same

factorizations of rs . If u is $r = d_1 \cdot d_2 \cdot \dots \cdot d_i$ and v is $s = e_1 \cdot e_2 \cdot \dots \cdot e_j$, we define the factorizations of rs

$$\begin{aligned} F((u, v)) &= d_1 \cdot d_2 \cdot \dots \cdot d_i \cdot e_1 \cdot e_2 \cdot \dots \cdot e_j \\ G((u, v)) &= d_1 \cdot d_2 \cdot \dots \cdot d_{i-1} \cdot (d_i e_1) \cdot e_2 \cdot \dots \cdot e_j. \end{aligned}$$

The inequality (6) follows from the fact that the mappings F and G are injections from X to Y which moreover have disjoint images. We leave a simple verification of this fact to the reader.

Suppose now that $m_k(n_0) > n_0^{\rho_k}/\sqrt{2}$ for some $n_0 \geq 2$. By (6), we have $m_k(n_0^2) \geq 2m_k(n_0)^2 > n_0^{2\rho_k}$ and hence we can take some $\varepsilon > 0$ so that $m_k(n_0^2) \geq (n_0^2)^{\rho_k+\varepsilon}$. Then, again by (6), $m_k(n_0^{2^i}) \geq (n_0^{2^i})^{\rho_k+\varepsilon}$ for every $i = 1, 2, \dots$, which is in contradiction with $m_k(n) = o(n^{\rho_k+\varepsilon})$. $\square$

Lemma 2.5. *Suppose that $q_1, \dots, q_k$ are primes, not necessarily distinct, such that the product $q_1 q_2 \dots q_k$ divides n . Then, with $m(1) = 1$,*

$$m(n) < (2\Omega(n))^k \cdot m(n/q_1 q_2 \dots q_k). \quad (7)$$

Proof. It suffices to prove only the case $k = 1$; i.e., the inequality

$$m(n) < 2\Omega(n) \cdot m(n/p), \quad (8)$$

where p is a prime dividing n , because the general case follows easily by iteration. Let X be the set of all pairs (u, i) where u is an ordered factorization of n/p (in parts bigger than 1) and i is an integer satisfying $1 \leq i \leq 2r + 1$, where r is the number of parts in u . Let Y be the set of all ordered factorizations of n in parts bigger than 1. We shall define a surjection F from X onto Y . This will prove (8) because $r \leq \Omega(n/p) = \Omega(n) - 1$, and therefore for every u we have $2r + 1 < 2\Omega(n)$ pairs (u, i) , and so

$$m(n) = |Y| \leq |X| < 2\Omega(n) \cdot m(n/p).$$

For $(u, i) \in X$, where u is $n/p = d_1 \cdot d_2 \cdot \dots \cdot d_r$, we define $j = i - r$ and set $F((u, i))$ to be the factorization

$$n = d_1 \cdot \dots \cdot d_{i-1} \cdot (pd_i) \cdot d_{i+1} \cdot \dots \cdot d_r$$

if $1 \leq i \leq r$ and

$$n = d_1 \cdot \dots \cdot d_{j-1} \cdot p \cdot d_j \cdot \dots \cdot d_r$$

if $r + 1 \leq i \leq 2r + 1$ (for $j = 1$, p is the first part and for $j = r + 1$ it is the last one). It is clear that F is a surjection. $\square$

Lemma 2.6. *If $n_1, n_2, \dots, n_k$ are positive integers such that for no $i \neq j$ we have $n_i | n_j$, then*

$$m(n_1 n_2 \dots n_k) \geq k! \cdot m(n_1) m(n_2) \dots m(n_k).$$

This implies that for every $n \geq 1$ we have

$$m(n) \geq \omega(n)! \cdot 2^{\Omega(n) - \omega(n)} \quad \text{and} \quad m(n) \geq 2^{\Omega(n) - 1}.$$

Proof. Let X be the set of all k -tuples $(u_1, u_2, \dots, u_k)$, where u_i is an ordered factorization of n_i in parts bigger than 1 and let Y be the set of these factorizations for $n_1 n_2 \dots n_k$. For every permutation σ of $1, 2, \dots, k$, we define a mapping $F_\sigma : X \rightarrow Y$ by

$$F_\sigma((u_1, u_2, \dots, u_k)) = u_{\sigma(1)} \cdot u_{\sigma(2)} \cdot \dots \cdot u_{\sigma(k)},$$

i.e., we concatenate factorizations u_i in the order prescribed by σ . It is clear that each F_σ is an injection. Suppose that $F_\sigma((u_1, u_2, \dots, u_k)) = F_\tau((v_1, v_2, \dots, v_k))$ for some permutations σ, τ and factorizations u_i and v_i . It follows that $u_{\sigma(1)}$ is an initial segment of $v_{\tau(1)}$ or vice versa, and hence $n_{\sigma(1)}$ divides $n_{\tau(1)}$ or vice versa. This implies that $\sigma(1) = \tau(1)$ and $u_{\sigma(1)} = v_{\tau(1)}$. Applying the same argument, we obtain that $\sigma(j) = \tau(j)$ and $u_{\sigma(j)} = v_{\tau(j)}$ also for $j = 2, \dots, k$. Thus $\sigma = \tau$ and $u_j = v_j$ for $j = 1, 2, \dots, k$. We have proved that the $k!$ mappings F_σ have mutually disjoint images. Therefore

$$k! \cdot m(n_1) m(n_2) \dots m(n_k) = k! |X| \leq |Y| = m(n_1 n_2 \dots n_k).$$

If $n = q_1^{a_1} q_2^{a_2} \dots q_k^{a_k}$ is the prime factorization of n , applying the first inequality to the k numbers $n_i = q_i^{a_i}$ and using that $m(p^a) = 2^{a-1}$, we obtain

$$m(n) \geq k! \prod_{i=1}^k 2^{a_i-1} = k! \cdot 2^{\Omega(n)-k},$$

which is the second inequality. Using that $k!/2^k \geq 1/2$ for every $k \geq 1$, we get the third inequality. $\square$

Note that $m(n) \geq 2^{\Omega(n)-1}$ is tight for every $n = p^a$.

3 The upper bound

We prove the following upper bound on the maximal order of $m(n)$.

Theorem 3.1. *For every $\varepsilon > 0$ we have*

$$m(n) < \frac{n^\rho}{\exp((\log n)^{1/\rho}/(\log \log n)^{1+\varepsilon+o(1)})}$$

for integers $n > 2$.

Proof. Let $\varepsilon > 0$ be given. To bound $m(n)$ from above, we split the integers $n > 0$ in two groups, those with $\omega(n) \leq k$ and those with $\omega(n) > k$, which we shall treat by different arguments; the optimum value of the parameter $k = k(n)$ will be selected in the end.

The case $\omega(n) \leq k$. Let $n = q_1^{a_1} q_2^{a_2} \dots q_r^{a_r}$, $r \leq k$, be the prime decomposition of n where $q_1 < q_2 < \dots < q_r$. We denote by $\bar{n}$ the number obtained from n by replacing q_i in the decomposition by p_i , the i th smallest prime. Then $\bar{n} \leq n$. From the fact that $m(n)$ depends only on the exponents a_i and from Lemma 2.4 we get

$$m(n) = m(\bar{n}) = m_r(\bar{n}) < \bar{n}^{\rho_r} \leq n^{\rho_k}.$$

Thus, by Lemma 2.3,

$$\begin{aligned} m(n) &< n^{\rho_k} \\ &= n^\rho \exp(-(\rho - \rho_k) \log n) \\ &= n^\rho \exp\left(-(c + o(1)) \frac{\log n}{k^{\rho-1} (\log k)^\rho}\right) \end{aligned} \tag{9}$$

where $c = (\rho - 1)^{-1} |\zeta'(\rho)|^{-1} > 0$.

The case $\omega(n) > k$. Let $l(n)$ be the product of some k distinct prime factors of n ; then $l(n) \geq p_1 p_2 \dots p_k$, the product of the k smallest primes. We have the estimates

$$\sum_{p \leq p_k} \log p = p_k + O(p_k / \log p_k) = k \log k + k \log \log k + O(k)$$

and

$$2\Omega(n) \leq (2/\log 2) \log n < 3 \log n.$$

By Lemmas 2.5, 2.4 and these estimates,

$$\begin{aligned}
m(n) &< (2\Omega(n))^k m(n/\ell(n)) < (3\log n)^k \frac{n^\rho}{\ell(n)^\rho} \\
&\leq (3\log n)^k \frac{n^\rho}{(p_1 \dots p_k)^\rho} \\
&= n^\rho \exp(-k(\rho \log k + \rho \log \log k - \log \log n + O(1))). \quad (10)
\end{aligned}$$

To determine the best upper bound on $m(n)$, we begin with k in the form $k = k(n) = (\log n)^{\alpha+o(1)}$ where $\alpha \in (0, 1)$ is a constant. Necessarily $\alpha \geq 1/\rho$, for else the argument of exp in (10) is eventually positive and we get a useless bound. It follows that the optimum is $\alpha = 1/\rho$ when the arguments of both exps in (9) and (10) are $-(\log n)^{1/\rho+o(1)}$, provided that

$$\rho \log k + \rho \log \log k - \log \log n + O(1) > c > 0 \quad (11)$$

for big n . Now we set, more precisely,

$$k = k(n) = \frac{(\log n)^{1/\rho}}{(\log \log n)^{d+o(1)}}$$

with a constant $d > 0$. With this k , the function in (11) becomes $\rho(1 - d + o(1)) \log \log \log n + O(1)$ and we see that condition (11) is satisfied for $d < 1$ (for $d > 1$ the argument of the exp in (10) is again eventually positive). With this k , the arguments of the exps in (9) and (10) are, respectively,

$$-\frac{(\log n)^{1/\rho}}{(\log \log n)^{1+(\rho-1)(1-d)+o(1)}} \quad \text{and} \quad -\frac{(\log n)^{1/\rho}}{(\log \log n)^{d+o(1)}}.$$

Setting $d = 1 - \varepsilon/2(\rho - 1)$, we obtain the stated bound. $\square$

4 The lower bound

We prove the following lower bound on the maximal order of $m(n)$.

Theorem 4.1. *There is a constant $c > 0$ such that the inequality*

$$m(n) > \frac{n^\rho}{\exp(c(\log n / \log \log n)^{1/\rho})}$$

holds for infinitely many integers $n > 0$.

We shall see that it is possible to take $c = 2.01630 \cdots - \varepsilon$. We begin with explaining the effective Ikehara–Ingham theorem on Dirichlet series. Then we apply it to $1/(2 - \zeta_k(s))$ to obtain an asymptotic relation for the average order of $m_k(n)$ with error estimate independent on k . Finally, combining this relation with an estimate on density of smooth numbers we obtain Theorem 4.1. For the background on Dirichlet series we refer to Tenenbaum [27].

Suppose that $(a_n)_{n \geq 1}$ is a sequence of nonnegative real numbers with the summatory function

$$A(t) = \sum_{n \leq e^t} a_n$$

and the Dirichlet series

$$F(s) = \sum_{n=1}^{\infty} \frac{a_n}{n^s} = \int_{0-}^{\infty} e^{-st} dA(t).$$

Suppose that $F(s)$ converges for $\sigma > a > 0$. We may assume that a is the abscissa of (absolute) convergence; then by the Phragmén–Landau theorem, a is a singularity of $F(s)$. The effective Ikehara–Ingham theorem, proved by Tenenbaum [27] (who used the method of Ganelius [5]), extracts an asymptotic relation for $A(x)$ as $x \rightarrow \infty$ from the local behavior of $F(s)$ near a and, moreover, it provides an explicit estimate of the error term in terms of the regularity of $F(s)$ on the vertical segments $a + \sigma + i\tau$, $-T \leq \tau \leq T$, as $\sigma \rightarrow 0+$. We quote the theorem verbatim from Tenenbaum [27, p. 234].

Theorem 4.2. (“Effective” Ikehara–Ingham). *Let $A(t)$ be a non-decreasing function such that the integral*

$$F(s) := \int_0^{\infty} e^{-st} dA(t)$$

converges for $\sigma > a > 0$. Suppose that there exist constants $c \geq 0$, $\omega > -1$, such that the function

$$G(s) := \frac{F(s+a)}{s+a} - \frac{c}{s^{\omega+1}} \quad (\sigma > 0)$$

satisfies

$$\eta(\sigma, T) := \sigma^{\omega} \int_{-T}^T |G(2\sigma + i\tau) - G(\sigma + i\tau)| d\tau = o(1) \quad (\sigma \rightarrow 0+) \quad (12)$$

for each fixed $T > 0$. Then we have

$$A(x) = \left\{ \frac{c}{\Gamma(\omega + 1)} + O(\rho(x)) \right\} e^{ax} x^\omega \quad (x \geq 1), \quad (13)$$

with

$$\rho(x) := \inf_{T \geq 32(a+1)} \{T^{-1} + \eta(1/x, T) + (Tx)^{-\omega-1}\}.$$

Furthermore, the implicit constant in (13) depends only on a , c , and ω . An admissible choice for this constant is

$$52 + 1652c(a+1)(\omega+1) + 69c(1 + (\omega+1)e^{1-\omega}(\omega+1)^{\omega+2})/\Gamma(\omega+1).$$

Note that for a meromorphic $F(s)$ with a simple pole at $s = a$ (so $\omega = 0$), the condition (12) is satisfied iff $F(s)$ has on the line $\sigma = a$ no other poles.

We shall apply Theorem 4.2 to the functions

$$F(s) = F_k(s) = \sum_{n \geq 1} \frac{m_k(n)}{n^s} = \frac{1}{2 - \zeta_k(s)}$$

for $k \geq 2$, $a = \rho_k$, $c = c_k = -1/\rho_k \zeta'_k(\rho_k)$, and $\omega = 0$. It is not hard to prove (we do this in the next Proposition) that ρ_k is the only pole of $F_k(s)$ on $\sigma = \rho_k$ when $k \geq 2$ (this is not true for $k = 1$) and thus by Theorem 4.2

$$\sum_{n \leq x} m_k(n) = (c_k + o(1))x^{\rho_k} \quad (x \rightarrow \infty)$$

for each fixed $k \geq 2$. (In contrast, $\sum_{n \leq x} m_1(n) = 2^r - 1$ where $2^r \leq x < 2^{r+1}$.) To get a good lower bound on $m(n)$, we have to strengthen this by obtaining uniformity in k of the error term $o(1)$. This follows from Theorem 4.2, once we prove that for $F(s) = F_k(s)$ the condition (12) is satisfied uniformly in k .

Proposition 4.3. *Let, for $k \geq 2$,*

$$G_k(s) = \frac{F_k(s + \rho_k)}{s + \rho_k} - \frac{c_k}{s} = \frac{1}{(2 - \zeta_k(s + \rho_k))(s + \rho_k)} - \frac{c_k}{s}$$

and $T > 0$ be arbitrary but fixed. Then

$$\lim_{\sigma \rightarrow 0+} \int_{-T}^T |G_k(2\sigma + i\tau) - G_k(\sigma + i\tau)| \, d\tau = 0$$

uniformly in $k \geq 2$; that is, the condition (12) holds uniformly in k .

Proof. Let $t(\sigma) = \sigma^{1/5}$; any function $t(\sigma) > 0$ satisfying, as $\sigma \rightarrow 0+$, that $t(\sigma) \rightarrow 0$ and $\sigma/t(\sigma)^4 \rightarrow 0$ would do in our argument. For every fixed $T > 0$, we bound the integrand by a quantity that depends only on σ and not on τ and $k \geq 2$ and that goes to 0 as $\sigma \rightarrow 0+$; this will prove the statement. We manage doing this by splitting $[-T, T]$ in two ranges, $t(\sigma) \leq |\tau| \leq T$ and $|\tau| \leq t(\sigma)$, in which we apply different arguments.

The range $t(\sigma) \leq |\tau| \leq T$. Denoting by γ the horizontal segment with endpoints $\sigma + i\tau$ and $2\sigma + i\tau$, we have the bound

$$|G_k(2\sigma + i\tau) - G_k(\sigma + i\tau)| = \left| \int_{\gamma} G'_k(z) dz \right| \leq \sigma |G'_k(s_0)|$$

where s_0 is some point lying on γ . The derivative of $G_k(s)$ equals

$$G'_k(s) = \frac{(s + \rho_k)\zeta'_k(s + \rho_k) + \zeta_k(s + \rho_k) - 2}{(2 - \zeta_k(s + \rho_k))^2(s + \rho_k)^2} + \frac{c_k}{s^2}.$$

We bound the numerators and denominators of this expression. As for the numerators, by Lemma 2.2, there is a constant $c = c(T) > 0$ depending only on T such that

$$|(s + \rho_k)\zeta'_k(s + \rho_k) + \zeta_k(s + \rho_k) - 2|, |c_k| < c$$

holds for every $k \geq 2$ and s with $0 < \sigma < 1$ and $|\tau| \leq T$. For the second denominator, we have, in our range and for $0 < \sigma < 1$,

$$\frac{\sigma}{|s_0|^2} \leq \frac{\sigma}{\sigma^2 + t(\sigma)^2} = \frac{\sigma^{3/5}}{\sigma^{8/5} + 1} < \sigma^{3/5}.$$

We bound the first denominator. Clearly, $|s + \rho_k|^2 \geq \rho_k^2 > 1$ for every s with $\sigma > 0$. For every $k \geq 2$ and every s with $0 < \sigma < 1$ and any τ we have

$$|2 - \zeta_k(s + \rho_k)| \geq \operatorname{Re}(2 - \zeta_k(s + \rho_k)) = \sum_{\substack{n \geq 1 \\ P(n) \leq p_k}} \frac{1}{n^{\rho_k + \sigma}} (n^{\sigma} - \cos(\tau \log n))$$

and, consequently, (recall that $k \geq 2$ and $1 < \rho_k < 2$)

$$|2 - \zeta_k(s + \rho_k)|^2 > \left(\frac{2 - \cos(\tau \log 2) - \cos(\tau \log 3)}{27} \right)^2 =: h(\tau).$$

Since $2^\alpha = 3$ holds for no fraction α , $h(\tau) = 0$ only for $\tau = 0$. The function $h(\tau)$ is continuous and even and $h(\tau) \sim \beta\tau^4$ as $\tau \rightarrow 0$ for a constant $\beta > 0$. Thus there is a constants $\beta_1 = \beta_1(T) < 1$ depending only on T such that if $0 < \sigma < \beta_1$ then the minimum of $h(\tau)$ on $[t(\sigma), T]$ is attained at $t(\sigma)$ and $h(t(\sigma)) > \beta t(\sigma)^4/2$. Hence, in our range and for $0 < 2\sigma < \beta_1$,

$$\frac{\sigma}{|2 - \zeta_k(s_0 + \rho_k)|^2 \cdot |s_0 + \rho_k|^2} < \frac{2\sigma}{\beta t(\sigma)^4} = \frac{2\sigma^{1/5}}{\beta}.$$

Taking together all estimates, we have in our range and for $0 < \sigma < \beta_1/2$ that

$$|G_k(2\sigma + i\tau) - G_k(\sigma + i\tau)| \leq \sigma |G'_k(s_0)| < c(2\sigma^{1/5}/\beta + \sigma^{3/5}),$$

which is the required bound.

The range $|\tau| \leq t(\sigma)$. We prove that there is an absolute constant $\delta > 0$ such that for every $k \geq 2$ and s with $|s| < \delta$ we have the expansion

$$G_k(s) = d_k + O(s),$$

where d_k is a constant and the constant implicit in O is absolute. (We need independence on k both for the constant in $O(s)$ and for the domain of validity of the error estimate.) Then if $0 < \sigma < \delta^5/32$ and $|\tau| \leq t(\sigma)$, both numbers $\sigma + i\tau$ and $2\sigma + i\tau$ satisfy $|s| < \delta$, and we have the bound

$$|G_k(2\sigma + i\tau) - G_k(\sigma + i\tau)| = O(|\sigma + i\tau| + |2\sigma + i\tau|) = O(\sigma^{1/5})$$

with absolute constants in O s, which is the required bound.

We begin with the origin-centered closed disc $B = B(0, 0.1)$; the point of the radius 0.1 is only that $\rho_2 - 0.1 > 1$. We define functions $f_k(s)$ by

$$f_k(s) = \frac{\zeta_k(s + \rho_k) - 2 - s\zeta'_k(\rho_k) - s^2\zeta''_k(\rho_k)/2}{s^3}.$$

Let a_k be the maximum value taken by $|\zeta_k(s)|$ on the circle $|s - \rho_k| = 0.1$. By the maximum modulus principle ($f_k(s)$ is holomorphic on B), for every $s \in B$ we have

$$|f_k(s)| \leq 10^3(a_k + 2 + 10^{-1}\zeta'_k(\rho_k) + 10^{-2}\zeta''_k(\rho_k)/2).$$

Thus, by Lemma 2.2, there is an absolute constant $M > 0$ such that

$$|f_k(s)| < M$$

holds for every $s \in B$ and every $k \geq 2$. We rewrite $\zeta_k(s + \rho_k) = 2 + s\zeta'_k(\rho_k) + s^2\zeta''_k(\rho_k)/2 + s^3f_k(s)$ as

$$\begin{aligned} \frac{1}{(2 - \zeta(s + \rho_k))(s + \rho_k)} &= -\frac{1}{s\rho_k\zeta'_k(\rho_k)} \times \frac{1}{1 + s/\rho_k} \\ &\times \frac{1}{1 + s\zeta''_k(\rho_k)/2\zeta'_k(\rho_k) + s^2f_k(s)/\zeta'_k(\rho_k)} \\ &= -\frac{1}{s\rho_k\zeta'_k(\rho_k)} \times \frac{1}{1 + s/\rho_k} \times \frac{1}{1 + sb_k + s^2h_k(s)}. \end{aligned}$$

It follows, by Lemma 2.2 and the bound $|f_k(s)| < M$ valid on B , that there is a δ , $0 < \delta < 0.1$, such that $|s/\rho_k| < 1/2$ and $|sb_k + s^2h_k(s)| < 1/2$ whenever $|s| < \delta$ and $k \geq 2$. Using the estimate $(1 + s)^{-1} = 1 - s + O(s^2)$, valid for $|s| < 1/2$, and Lemma 2.2 we obtain for $k \geq 2$ and $|s| < \delta$ the expansion

$$\begin{aligned} \frac{1}{(2 - \zeta_k(s + \rho_k))(s + \rho_k)} &= \frac{c_k}{s} \left(1 - \frac{s}{\rho_k} + O(s^2)\right) \left(1 - s \frac{\zeta''_k(\rho_k)}{2\zeta'_k(\rho_k)} + O(s^2)\right) \\ &= \frac{c_k}{s} - c_k \left(\frac{1}{\rho_k} + \frac{\zeta''_k(\rho_k)}{2\zeta'_k(\rho_k)}\right) + O(s), \end{aligned}$$

where $c_k = -1/\rho_k\zeta'_k(\rho_k)$ and the constants in O s are absolute. Now the required expansion $G_k(s) = d_k + O(s)$ (valid for $|s| < \delta$ and with an absolute constant in O) is immediate. $\square$

Corollary 4.4. *There is a constant $\beta_2 > 2$ such that for every $x > \beta_2$ and every $k \geq 2$ we have*

$$\sum_{\substack{n \leq x \\ P(n) \leq p_k}} m(n) = \sum_{n \leq x} m_k(n) > x^{\rho_k}/5.$$

Proof. By Theorem 4.2 and Proposition 4.3, there is a function $e(x) > 0$ such that $e(x) \rightarrow 0$ as $x \rightarrow \infty$, and for every $x \geq 1$ and every $k \geq 2$ we have

$$\left| \sum_{n \leq x} m_k(n) - c_k x^{\rho_k} \right| < e(x) x^{\rho_k}.$$

The sequence of $c_k = -1/\rho_k\zeta'_k(\rho_k)$, $k = 1, 2, \dots$, monotonically decreases and converges to $c_\infty = -1/\rho\zeta'(\rho) > 0.3$. Thus if x is big enough so that $e(x) < 0.1$, the sum $\sum_{n \leq x} m_k(n)$ must be bigger than $0.2x^{\rho_k}$. $\square$

We proceed to the proof of Theorem 4.1. We denote, as usual,

$$\Psi(x, y) = \#\{n \leq x : P(n) \leq y\}.$$

By Corollary 4.4, for every $k \geq 2$ and $x > \beta_2$ there exists an $n_0 \leq x$ such that

$$\Psi(x, p_k)m(n_0) > x^{\rho_k}/5 = \frac{x^\rho}{5 \exp((\rho - \rho_k) \log x)}.$$

We select $k = k(x)$ so that it satisfies

$$k = (\log x)^{\alpha+o(1)}$$

as $x \rightarrow \infty$, for some absolute constant $\alpha \in (0, 1)$ (we make our choice of k more precise later). Then

$$p_k = (1 + o(1))k \log k = (\log x)^{\alpha+o(1)}.$$

A theorem due to de Bruijn, see Theorem 2 in Tenenbaum's book [27, p. 359], shows that

$$\log(\Psi(x, p_k)) = (1 + o(1))Z,$$

where

$$\begin{aligned} Z &= \frac{\log x}{\log p_k} \log \left(1 + \frac{p_k}{\log x} \right) + \frac{p_k}{\log p_k} \log \left(1 + \frac{\log x}{p_k} \right) \\ &= \frac{p_k}{\log p_k} (1 + o(1)) + \frac{p_k}{\log p_k} \log \left(1 + \frac{\log x}{p_k} \right) \\ &= (1 + o(1))k(\log \log x - \log k). \end{aligned}$$

By Lemma 2.3,

$$\rho - \rho_k = \frac{c_1 + o(1)}{k^{\rho-1}(\log k)^\rho}$$

where $c_1 = 1/((\rho-1)|\zeta'(\rho)|)$. Substituting both estimates in the lower bound on $\Psi(x, p_k)m(n_0)$, we get (absorbing the 5 in the denominator in the $o(1)$ terms)

$$m(n_0) > \frac{x^\rho}{\exp \left(c_1(1 + o(1)) \frac{\log x}{k^{\rho-1}(\log k)^\rho} + (1 + o(1))k(\log \log x - \log k) \right)}.$$

This suggests to choose k so that both terms in the argument of the exponential,

$$\frac{\log x}{k^{\rho-1}(\log k)^\rho} \quad \text{and} \quad k(\log \log x - \log k),$$

are of the same order of magnitude. This occurs when $\alpha = 1/\rho$, more precisely when

$$k = \lfloor d(\log x)^{1/\rho}(\log \log x)^{-(\rho+1)/\rho} \rfloor$$

with any constant $d > 0$, because then

$$\frac{\log x}{k^{\rho-1}(\log k)^\rho} = d^{1-\rho} \rho^\rho (1 + o(1)) \left(\frac{\log x}{\log \log x} \right)^{1/\rho}$$

and

$$k(\log \log x - \log k) = (1 - \rho^{-1})d(1 + o(1)) \left(\frac{\log x}{\log \log x} \right)^{1/\rho}.$$

Thus, for this selection of k ,

$$m(n_0) > \frac{x^\rho}{\exp \left((c + o(1)) \left(\frac{\log x}{\log \log x} \right)^{1/\rho} \right)}$$

where $c > 0$ is a constant depending only on the choice of d . The lower bound eventually increases monotonically to infinity, and we conclude that there exist infinitely many numbers n_0 satisfying

$$m(n_0) > \frac{n_0^\rho}{\exp \left((c + o(1)) \left(\frac{\log n_0}{\log \log n_0} \right)^{1/\rho} \right)}.$$

The proof of Theorem 4.1 is complete.

It is not difficult to find the optimal value of d ; it yields the value

$$c = (\rho^{\rho+1} c_1)^{1/\rho} = \left(\frac{\rho^{\rho+1}}{(\rho-1)|\zeta'(\rho)|} \right)^{1/\rho} \approx 2.01630.$$

5 Historical remarks and arithmetical properties of $m(n)$

We begin with a survey of some previous results on $m(n)$. We restrict our attention only to works dealing directly with this quantity. There are many other variants of factorization counting functions (with restrictions on factors, counting unordered factorizations etc.) and for a survey on these we refer the reader to Knopfmacher and Mays [16].

Kalmár proved in [14] that the error term $o(1)$ in (1) is

$$O(\exp(-\alpha \log \log x \cdot \log \log \log x)), \text{ with } \alpha < \frac{1}{2(\rho - 1) \log 2} \approx 1.97996.$$

Ikehara devoted three papers to the estimates of $M(x)$. In [10], he gave weak bounds of the type $M(x) > x^{\rho-\varepsilon}$ on a sequence of x tending to infinity, and $M(x) < x^{\rho+\varepsilon}$ for all large enough x . In the review of [10], Kalmár pointed out a gap in the proof and sketched a correct argument. In [11], Ikehara gave a proof of (1) with an error bound $O(\exp(q \log \log x))$ for some constant $q < 0$, which is slightly weaker than Kalmár's result. Finally, in [12], he succeeded to get a stronger error bound

$$O(\exp(-\alpha(\log \log x)^\gamma)), \text{ with } \alpha > 0 \text{ and } \gamma < 4/3.$$

Hwang [9] obtained an improvement of Ikehara's last bound by replacing $4/3$ with $3/2$.

Rieger proved in [23], besides other results, that for all positive integers k, l with $(k, l) = 1$ one has

$$\sum_{n \leq x, n \equiv l \pmod{k}} m(n) = \frac{1 + o(1)}{\varphi(k)} M(x) = \frac{-1}{\varphi(k) \rho \zeta'(\rho)} \cdot x^\rho (1 + o(1)).$$

Warlimont investigated in [28] variants of $m(n)$ counting ordered factorizations with distinct parts and with coprime parts and estimated their summatory functions. Hille in [8] proved that $m(n) = O(n^\rho)$ and that $m(n) > n^{\rho-\varepsilon}$ for infinitely many n . We already mentioned in Section 1 the remark of Erdős on $m(n)$ in [4] and we mentioned (and improved) the result of Chor, Lemke and Mador [1] that $m(n) < n^\rho$ for all n . Other elementary and constructive

proofs of the bounds $m(n) \leq n^\rho$ and $\limsup_n m(n)/n^{\rho-\epsilon} = \infty$ were recently given by Coppersmith and Lewenstein [3].

We now turn to recurrences and explicit formulas. The recurrence $m(1) = 1$ and

$$m(n) = \sum_{d|n, d < n} m(d) \quad \text{for } n > 1 \quad (14)$$

is immediate from fixing the first part in a factorization. If we set $m^*(1) = 1/2$ and $m^*(n) = m(n)$ for $n > 1$, then $2m^*(n) = \sum_{d|n} m(d)$ holds for all $n \geq 1$. By Möbius inversion, $m(n) = 2 \sum_{d|n} \mu(d) m^*(n/d)$ for all $n \geq 1$. For $n = q_1^{a_1} q_2^{a_2} \dots q_r^{a_r} > 1$ this can be rewritten as the recurrence formula

$$m(n) = 2 \left(\sum_i m\left(\frac{n}{q_i}\right) - \sum_{i < j} m\left(\frac{n}{q_i q_j}\right) + \dots + (-1)^{r-1} m\left(\frac{n}{q_1 q_2 \dots q_r}\right) \right), \quad (15)$$

in which we must set $m(1) = 1/2$. Formulas (14) and (15) are from Hille's paper [8]. In fact, (15) is stated there incorrectly with $m(1) = 1$, as was pointed out by Kühnel [17] and Sen [24].

Clearly, $m(p^a) = 2^{a-1}$ because ordered factorizations of p^a in parts > 1 are in bijection with (additive) compositions of a in parts > 0 . If $p \neq q$ are primes and $a \geq b \geq 0$ are integers, we have the formula

$$m(p^a q^b) = 2^{a+b-1} \sum_{k=0}^b \binom{a}{k} \binom{b}{k} 2^{-k}$$

that was derived in [1] and before by Sen [24] and MacMahon [21]. In particular,

$$m(p^a q) = (a+2)2^{a-1} \quad \text{and} \quad m(p^a q^2) = (a^2 + 7a + 8)2^{a-2}. \quad (16)$$

In general, for $n = q_1^{a_1} q_2^{a_2} \dots q_r^{a_r}$, and $a = a_1 + a_2 + \dots + a_r$, MacMahon [21] derived the formula

$$m(q_1^{a_1} q_2^{a_2} \dots q_r^{a_r}) = \sum_{j=1}^a \sum_{i=0}^{j-1} (-1)^i \binom{j}{i} \prod_{k=1}^r \binom{a_k + j - i - 1}{a_k}.$$

A more complicated summation formula for $m(q_1^{a_1} q_2^{a_2} \dots q_r^{a_r})$ but involving only nonnegative summands was obtained by Kühnel in [17] and [18]. Let

$d_k(n)$ be the number of solutions of $n = n_1 n_2 \dots n_k$, where $n_i \geq 1$ are positive integers; so $d_2(n)$ is the number of divisors of n . Sklar [25] mentions the formula

$$m(n) = \sum_{k=1}^{\infty} \frac{d_k(n)}{2^{k+1}}. \quad (17)$$

Somewhat surprisingly, $m(n)$ has an additive definition in terms of integer partitions. We say that a partition $(1^{a_1}, 2^{a_2}, \dots, k^{a_k})$ of n is *perfect*, if for every $m < n$ there is exactly one k -tuple $(b_1, \dots, b_k)$, $0 \leq b_i \leq a_i$ for all i , such that $(1^{b_1}, 2^{b_2}, \dots, k^{b_k})$ is a partition of m . MacMahon [19] proved the identity

$$m(n) = \# \text{ perfect partitions of } (n-1).$$

For example, since $m(12) = 8$, we have 8 perfect partitions of 11, namely $(1^2, 3, 6)$, $(1, 2^2, 6)$, $(1^5, 6)$, $(1, 2, 4^2)$, $(1^3, 4^2)$, $(1^2, 3^3)$, $(1, 2^5)$, and (1^{11}) .

In conclusion of the survey of previous results we should remark that from an enumerative point of view it is natural to consider $m(n)$ as a function of the partition $\lambda = (a_1, a_2, \dots, a_k)$ of $\Omega(n)$, where $n = q_1^{a_1} q_2^{a_2} \dots q_k^{a_k}$ with $a_1 \geq a_2 \geq \dots \geq a_k$, rather than n . Then $m(\lambda)$ is defined as the number of ways to write $\lambda = v_1 + v_2 + \dots + v_t$ where each v_i is a k -tuple of nonnegative integers, the order of summands matters, and no v_i is a zero vector. So $m(\lambda)$ is naturally understood as the number of k -dimensional compositions of λ . This approach was pursued by MacMahon in his memoirs [19], [20], and [21], see also [22].

The sequence

$$(m(n))_{n \geq 1} = (1, 1, 1, 2, 1, 3, 1, 4, 2, 3, 1, 8, 1, 3, 3, 8, 1, 8, 1, 8, 3, 3, 1, 20, 2, \dots)$$

forms entry A074206 of the database [26]. Continuing the sequence a little further, we notice that $m(48) = 48$ and that $n = 48 = 2^4 \cdot 3$ is the smallest $n > 1$ such that $m(n) = n$. The first formula in (16) produces infinitely many n with this property: setting $n = 2^{2q-2}q$ with a prime $q > 2$, we get $m(n) = n$. We record this observation as follows:

Proposition 5.1. *There exist infinitely many positive integers n such that $m(n) = n$.*

This result was obtained independently also by Knopfmacher and Mays [16].

We look at periodicity properties of the numbers $m(n)$. The recurrence (15) implies easily the following result.

Proposition 5.2. *The number $m(n)$ is odd if and only if n is squarefree.*

It would be interesting to characterize the behavior of $m(n)$ with respect to other moduli besides 2. In the next Proposition we give a partial result in this direction. Recall that an integer valued function $f(n)$ defined on the set of positive integers is called *eventually periodic* modulo k if there exist integers n_0 and T such that $f(n) \equiv f(n+T) \pmod{k}$ for all $n > n_0$. We show that $m(n)$ is not eventually periodic modulo k by proving a stronger result that $m(n)$ is not eventually constant modulo k on any infinite arithmetic progression with coprime difference and the first term.

Proposition 5.3. *The function $m(n)$ is not eventually constant modulo k , where $k \geq 2$, on any infinite arithmetic progression $n \equiv A \pmod{K}$, $K \geq 2$, with coprime A and K .*

Proof. By Dirichlet's theorem, this arithmetic progression contains infinitely many prime numbers and therefore $m(n) = 1$ for infinitely many $n \equiv A \pmod{K}$. We select a prime q not dividing K and an integer z (coprime with K) such that $qz \equiv A \pmod{K}$. Since there are infinitely many prime numbers congruent to z modulo K , there are also infinitely many $n \equiv A \pmod{K}$ of the form qp where p is a prime. Thus there are infinitely many $n \equiv A \pmod{K}$ with $m(n) = 3$. Because $1 \not\equiv 3 \pmod{k}$ for $k > 2$, we are done if $k > 2$. For $k = 2$, $m(n) \equiv 1 \pmod{2}$ for infinitely many $n \equiv A \pmod{K}$ as before. As we noted, $m(n)$ is even iff n is not squarefree. It follows that $m(n) \equiv 0 \pmod{2}$ for infinitely many $n \equiv A \pmod{K}$ as well, which settles the case $k = 2$. $\square$

The condition $(A, K) = 1$ cannot be omitted because if (A, K) is not square-free, $m(n)$ is even for all $n \equiv A \pmod{K}$.

Recall now that a sequence $(f(n))_{n \geq 1}$ is holonomic if there exist positive integer polynomials $g_0, \dots, g_k$, not all zero, such that

$$g_k(n)f(n+k) + g_{k-1}(n)f(n+k-1) + \dots + g_0(n)f(n) = 0 \quad \text{for all } n \geq 1. \quad (18)$$

Proposition 5.4. *The sequence $m(n)$ is not holonomic.*

Proof. Dividing (18) by one of the (nonzero) coefficients g_j with the largest degree, we obtain the relation

$$f(n+j) = \sum_{0 \leq i \leq k, i \neq j} h_i(n)f(n+i)$$

where the h_i 's are rational functions such that each $h_i(x)$ goes to a finite constant c_i as $x \rightarrow \infty$ (we may even assume that $|c_i| \leq 1$ for every i). Hence there is a constant $C > 0$ (depending only on k and the polynomials g_i) such that

$$|f(n)| \leq C \max \{|f(n+i)| : -k \leq i \leq k, i \neq 0\} \quad \text{for every } n \geq k+1.$$

We show that $(m(n))_{n \geq 1}$ violates this property.

We fix two integers $k, a \geq 1$ with the only restriction that a is coprime to each of the numbers $1, 2, \dots, k$. It is an easy consequence of the Fundamental Lemma of the Combinatorial Sieve (see [6]) that there is a constant $K > 0$ depending only on k so that

$$\Omega((an - k)(an - k + 1) \dots (an - 1)(an + 1) \dots (an + k)) \leq K$$

holds for infinitely many integers $n \geq 1$. For each of these n 's the $2k$ values $m(an+i)$, $-k \leq i \leq k$ and $i \neq 0$, are bounded by a constant (depending only on k) while the value $m(an)$ is at least $m(a)$ and can be made arbitrarily large by an appropriate selection of a . This contradicts the above property of holonomic sequences. $\square$

Remark 5.5. *The above proof can be adapted in a straightforward way to show that other number theoretical functions such as $\omega(n)$, $\Omega(n)$ and $\tau(n)$, where $\tau(n)$ is the number of divisors of n , are not holonomic.*

We present two more estimates related to the function $m(n)$.

Proposition 5.6. *The estimate*

$$\#\{m(n) : n \leq x\} \leq \exp\left(\pi\sqrt{2/\log 8}(1+o(1))(\log x)^{1/2}\right)$$

holds as $x \rightarrow \infty$.

Proof. Because $m(n)$ depends only on the partition $a_1 + \dots + a_k = \Omega(n)$, where $n = q_1^{a_1} \dots q_k^{a_k}$ ($q_1, \dots, q_k$ are distinct primes and $a_1 \geq a_2 \geq \dots \geq a_k > 0$ are integers), we have that

$$\#\{m(n) : n \leq x\} \leq p(1) + p(2) + \dots + p(r) \leq rp(r)$$

where $p(n)$ denotes the number of partitions of n and $r = \max_{n \leq x} \Omega(n)$. The result follows from $r \leq \log x / \log 2$ and the classic asymptotic relation $p(n) \sim \exp(\pi\sqrt{2n/3})/(4n\sqrt{3})$ due to Hardy and Ramanujan [7]. $\square$

We show that the same bound on the number of distinct values of $m(n)$ holds when the condition $n \leq x$ is replaced with $m(n) \leq x$.

Proposition 5.7. *The estimate*

$$\#\{m(n) : m(n) \leq x, n \geq 1\} \leq \exp\left(\pi\sqrt{2/\log 8}(1+o(1))(\log x)^{1/2}\right)$$

holds as $x \rightarrow \infty$.

Proof. As in Proposition 5.6 we have

$$\#\{m(n) : m(n) \leq x, n \geq 1\} \leq p(1) + p(2) + \cdots + p(r) \leq rp(r)$$

where now $r = \max_{m(n) \leq x} \Omega(n)$. By the third inequality in Lemma 2.6, $2^{r-1} = 2^{\Omega(n)-1} \leq m(n) \leq x$ for some n . Thus $r \leq 1 + \log x / \log 2$ and the result follows as in the proof of Proposition 5.6 using the asymptotics of $p(n)$. $\square$

References

- [1] B. Chor, P. Lemke and Z. Mador, ‘On the number of ordered factorizations of natural numbers’, *Discrete Math.* **214** (2000), 123–133.
- [2] M. Cipolla, ‘La determinazione assintotica dell’ n^{imo} numero primo’, *Napoli Rend.* (3) **8** 1902, 132–166.
- [3] D. Coppersmith and M. Lewenstein, ‘Constructive bounds on ordered factorizations’, *SIAM J. Discrete Math.* **19** (2005), 301–303.
- [4] P. Erdős, ‘On some asymptotic formulas in the theory of “factorisationum”’, *Ann. of Math. (2)* **42** (1941), 989–993.
- [5] T. Ganelius, *Tauberian Remainder Theorems*, Lecture Notes in Mathematics 232, Springer, Berlin, 1971.
- [6] H. Halberstam and H.-E. Richert, *Sieve Methods*, Academic Press, London, 1974.
- [7] G. H. Hardy and S. Ramanujan, ‘Asymptotic formulae in combinatory analysis’, *Proc. London Math. Soc. (2)* **17** (1918), 75–115.

- [8] E. Hille, ‘A problem in ”factorisatio numerorum”’, *Acta Arith.* **2** (1937), 134–146.
- [9] H.-K. Hwang, ‘Distribution of the number of factors in random ordered factorizations of integers’, *J. Number Theory* **81** (2000), 61–92.
- [10] S. Ikehara, ‘A theorem in ”factorisatio numerorum”’, *Tohoku Math. J.* **44** (1937), 162–164.
- [11] S. Ikehara, ‘On Kalmár’s problem in ”factorisatio numerorum”’, *Proc. Physic.-Math. Soc. Japan* **(3) 21** (1939), 208–219.
- [12] S. Ikehara, ‘On Kalmár’s problem in ”factorisatio numerorum.” II’, *Proc. Physic.-Math. Soc. Japan* **(3) 23** (1941), 767–774.
- [13] L. Kalmár, ‘A ”factorisatio numerorum” problémájáról’, *Matematikai és Fizikai Lapok* **38** (1931), 1–15.
- [14] L. Kalmár, ‘Über die mittlere Anzahl der Produktdarstellungen der Zahlen. (Erste Mitteilung)’, *Acta Litt. Sci. Szeged* **5** (1931), 95–107.
- [15] M. Klazar and F. Luca, ‘On the maximum order of numbers in the “factorisatio numerorum” problem’, preprint math.NT/0505352 in ArXiv, <http://www.arxiv.org/abs/math.NT/0505352>.
- [16] A. Knopfmacher and M. E. Mays, ‘A survey of factorization counting functions’, *Int. J. Number Theory* **1** (2005), 563–581.
- [17] U. Kühnel, *Über die Anzahl der Produktdarstellungen der positiven ganzen Zahlen*, dissertation (unpublished), 1941, 26 pages.
- [18] U. Kühnel, ‘Über die Anzahl der Produktdarstellungen der positiven ganzen Zahlen’, *Arch. Math.* **2** (1950), 216–219.
- [19] P. A. MacMahon, ‘The theory of perfect partitions of numbers and the compositions of multipartite numbers’, *Messenger Math.* **20** (1891), 103–119.
- [20] P. A. MacMahon, Yoke-chains and multipartite compositions in connexion with the analytical forms called trees, *Proc. London Math. Soc.* **22** (1891), 330–346.

- [21] P. A. MacMahon, ‘Memoir on the theory of the compositions of numbers’, *Philos. Trans. Roy. Soc. London (A)* **184** (1893), 835–901.
- [22] P. A. MacMahon, ‘Combinatory Analysis’, Cambridge University Press, 1915 and 1916 (Volume 1 and 2). Reprinted in one volume by Chelsea, 1960.
- [23] G. J. Rieger, ‘Über die Anzahl der Produktzerlegungen ganzer Zahlen’, *Math. Z.* **76** (1961), 226–234.
- [24] D. N. Sen, ‘A problem on ”factorisatio numerorum”’, *Bull. Calcutta Math. Soc.* **33** (1941), 1–8.
- [25] A. Sklar, ‘On the factorization of squarefree integers’, *Proc. Amer. Math. Soc.* **3** (1952), 701–705.
- [26] N. J. A. Sloane (ed.), The on-line encyclopedia of integer sequences, 2002, published electronically at <http://www.research.att.com/~njas/sequences>
- [27] G. Tenenbaum, *Introduction to analytic and probabilistic number theory*, Cambridge University Press, Cambridge 1995.
- [28] R. Warlimont, ‘Factorisatio numerorum with constraints’, *J. Number Theory* **45** (1993), 186–199.